

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

to forbid all communications. It is to be expected that with the precautionary measures that are being taken the disease will be stamped out before it assumes an epidemic form. One death from an infectious fever, probably enteric fever, has also occurred in this city, as will be seen by the mortuary statistics. I have been assured that due sanitary measures have been taken, as heretofore with all contagious and infectious diseases to prevent further issues.

Mortuary statistics for the last ten days of January, 1904.

Causes of death.	Number of deaths.	
Tuberculosis Septicæmia Infectious fever Athrepsia Lesion of heart, organic Congestion, cerebral Enteritis Bronchitis, capillary Debility, congenital Enteritis (under 2 years of age) Broncho-pneumonia	7 1 1 2 1 1 1 1 1	27 20 65 105 79 64 106 90 157 105
Total	18	

Estimated population, 48,000; annual rate of mortality per thousand, 13.68. No quarantinable disease has been reported.

Report from Santiago—Disinfecting barge San Rafael made ready for service.

Acting Assistant Surgeon Wilson reports, February 2, as follows: During the week ended January 30, 1904, bills of health were issued to three vessels bound for the United States.

No quarantinable disease has been reported.

The port quarantine service is better prepared for its duties now than it has been for some time, as the disinfecting barge San Rafael has been anchored in the bay, being in good working order, and it has received a new boiler for the steam launch which will be ready for use in a few days.

Mortality report for the week ended January 30, 1904.

Causes of death.	Number.	Bertillon number.
Tubercle of lungs. Syphilis (hereditary) Sarcoma Epithelioma Tetanus, infantile Arterio-sclerosis Pleuro-pneumonia Fibrinous pneumonia Intestinal obstruction Total.	5 1 1 1 1 1 1 1 1 1 1	27 36 45 45 72 81 93 93 108

Annual rate of mortality for the week, 14.85 per 1,000. Estimated population, 45,500.